

# Developmental changes in sensory-evoked optical intrinsic signals in the rat barrel cortex

Sintsov M., Suchkov D., Khazipov R., Minlebaev M.  
*Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia*

---

## Abstract

© 2017 Sintsov, Suchkov, Khazipov and Minlebaev. Optical Intrinsic Signal imaging (OISi) is a powerful technique for optical brain studies. OIS mainly reflects the hemodynamic response (HR) and metabolism, but it may also involve changes in tissue light scattering (LS) caused by transient cellular swelling in the active tissue. Here, we explored the developmental features of sensory-evoked OIS in the rat barrel cortex during the first 3 months after birth. Multispectral OISi revealed that two temporally distinct components contribute to the neonatal OIS: an early phase of LS followed by a late phase of HR. The contribution of LS to the early response was also evidenced by an increase in light transmission through the active barrel. The early OIS phase correlated in time and amplitude with the sensory-evoked electrophysiological response. Application of the Modified Beer-Lambert Law (MBLL) to the OIS data revealed that HR during the early phase involved only a slight decrease in blood oxygenation without any change in blood volume. In contrast, HR during the late phase manifested an adult-like increase in blood volume and oxygenation. During development, the peak time of the delayed HR progressively shortened with age, nearly reaching the stimulus onset and overlapping with the early LS phase by the fourth postnatal week. Thus, LS contributes to the sensory-evoked OIS in the barrel cortex of rats at all ages, and it dominates the early OIS phase in neonatal rats due to delayed HR. Our results are also consistent with the delayed blood oxygen level dependent (BOLD) signal in human preterm infants.

<http://dx.doi.org/10.3389/fncel.2017.00392>

---

## Keywords

Barrel field, Development, Hemodynamic response, Light scattering, Neonatal, OIS, Rats, Transparency

## References

- [1] Aitken P. G., Fayuk D., Somjen G. G., Turner D. A. (1999). Use of intrinsic optical signals to monitor physiological changes in brain tissue slices. *Methods* 18, 91–103. DOI: 10.1006/meth.1999.0762
- [2] Allievi A. G., Arichi T., Tusor N., Kimpton J., Arulkumaran S., Counsell S. J., et al. (2016). Maturation of sensorimotor functional responses in the preterm brain. *Cereb. Cortex* 26, 402–413. DOI: 10.1093/cercor/bhv203
- [3] Arichi T., Fagiolo G., Varela M., Melendez-Calderon A., Allievi A., Merchant N., et al. (2012). Development of BOLD signal hemodynamic responses in the human brain. *Neuroimage* 63, 663–673. DOI: 10.1016/j.neuroimage.2012.06.054
- [4] Attwell D., Buchan A. M., Charkpak S., Lauritzen M., Macvicar B. A., Newman E. A. (2010). Glial and neuronal control of brain blood flow. *Nature* 468, 232–243. DOI: 10.1038/nature09613

- [5] Bonhoeffer T., Grinvald A. (1991). Iso-orientation domains in cat visual cortex are arranged in pinwheel-like patterns. *Nature* 353, 429–431.
- [6] Borgdorff A. J., Poulet J. F. A., Petersen C. C. H. (2007). Facilitating sensory responses in developing mouse somatosensory barrel cortex. *J. Neurophysiol.* 97, 2992–3003. DOI: 10.1152/jn.00013.2007
- [7] Born A. P., Miranda M. J., Rostrup E., Toft P. B., Peitersen B., Larsson H. B., et al. (2000). Functional magnetic resonance imaging of the normal and abnormal visual system in early life. *Neuropediatrics* 31, 24–32. DOI: 10.1055/s-2000-15402
- [8] Buxton R. B., Uluda K., Dubowitz D. J., Liu T. T. (2004). Modeling the hemodynamic response to brain activation. *Neuroimage* 23(Suppl. 1), 220–233. DOI: 10.1016/j.neuroimage.2004.07.013
- [9] Colonnese M. T., Kaminska A., Minlebaev M., Milh M., Bloem B., Lescure S., et al. (2010). A conserved switch in sensory processing prepares developing neocortex for vision. *Neuron* 67, 480–498. DOI: 10.1016/j.neuron.2010.07.015
- [10] Cui X., Bray S., Bryant D. M., Glover G. H., Reiss A. L. (2011). A quantitative comparison of NIRS and fMRI across multiple cognitive tasks. *Neuroimage* 54, 2808–2821. DOI: 10.1016/j.neuroimage.2010.10.069
- [11] Daw M. I., Ashby M. C., Isaac J. T. R. (2007). Coordinated developmental recruitment of latent fast spiking interneurons in layer IV barrel cortex. *Nat. Neurosci.* 10, 453–461. DOI: 10.1038/nn1866
- [12] Delpy D. T., Cope M., Van Der Zee P., Arridge S., Wrayt S., Wyatt J., et al. (1988). Estimation of optical pathlength through tissue from direct time of flight measurement. *Phys. Med. Biol.* 33, 1433–1442. DOI: 10.1088/0031-9155/33/12/008
- [13] Fang Q., Boas D. A. (2009). Monte carlo simulation of photon migration in 3D turbid media accelerated by graphics processing units. *Opt. Exp.* 17:20178. DOI: 10.1364/OE.17.020178
- [14] Greisen G. (2005). Autoregulation of cerebral blood flow in newborn babies. *Early Hum. Dev.* 81, 423–428. DOI: 10.1016/j.earlhumdev.2005.03.005
- [15] Greisen G., Leung T., Wolf M. (2011). Has the time come to use near-infrared spectroscopy as a routine clinical tool in preterm infants undergoing intensive care? *Philos. Trans. Ser. A Maht. Phys. Eng. Scie.* 369, 4440–4451. DOI: 10.1098/rsta.2011.0261
- [16] Grinvald A., Lieke E., Frostig R. D., Gilbert C. D., Wiesel T. N. (1986). Functional architecture of cortex revealed by optical imaging of intrinsic signals. *Nature* 324, 361–364.
- [17] Grubb R. L., Raichle M. E., Eichling J. O., Ter-Pogossian M. M. (1974). The effects of changes in PaCO<sub>2</sub> cerebral blood volume, blood flow, and vascular mean transit time. *Stroke* 5, 630–639. DOI: 10.1161/01.STR.5.5.630
- [18] Hall C. N., Reynell C., Gesslein B., Hamilton N. B., Mishra A., Sutherland B. A., et al. (2014). Capillary pericytes regulate cerebral blood flow in health and disease. *Nature* 508, 55–60. DOI: 10.1038/nature13165
- [19] Harris J. J., Reynell C., Attwell D. (2011). The physiology of developmental changes in BOLD functional imaging signals. *Dev. Cogn. Neurosci.* 1, 199–216. DOI: 10.1016/j.dcn.2011.04.001
- [20] Hill R. A., Tong L., Yuan P., Murikinati S., Gupta S., Grutzendler J. (2015). Regional blood flow in the normal and ischemic brain is controlled by arteriolar smooth muscle cell contractility and not by capillary pericytes. *Neuron* 87, 95–110. DOI: 10.1016/j.neuron.2015.06.001
- [21] Hillman E. M. C. (2008). Optical brain imaging in vivo: techniques and applications from animal to man. *J. Biomed. Opt.* 12:051402. DOI: 10.1117/1.2789693
- [22] Huppert T., Hoge R., Diamond S., Franceschini M., Boas D. (2006). A temporal comparison of BOLD, ASL, and NIRS hemodynamic responses to motor stimuli in adult humans. *Neuroimage* 29, 368–382. DOI: 10.1016/j.neuroimage.2005.08.065
- [23] Jobsis F. (1977). Noninvasive, infrared monitoring of cerebral and myocardial oxygen sufficiency and circulatory parameters. *Science* 198, 1264–1267.
- [24] Jolivet R., Coggan J. S., Allaman I., Magistretti P. J. (2015). Multi-timescale modeling of activity-dependent metabolic coupling in the neuron-glia-vasculature ensemble. *PLoS Comput. Biol.* 11:e1004036. DOI: 10.1371/journal.pcbi.1004036
- [25] Jones M., Berwick J., Johnston D., Mayhew J. (2001). Concurrent optical imaging spectroscopy and laser-doppler flowmetry: the relationship between blood flow, oxygenation, and volume in rodent barrel cortex. *Neuroimage* 13, 1002–1015. DOI: 10.1006/nimg.2001.0808
- [26] Kalatsky V. A., Polley D. B., Merzenich M. M., Schreiner C. E., Stryker M. P. (2005). Fine functional organization of auditory cortex revealed by Fourier optical imaging. *Proc. Natl. Acad. Sci. U.S.A.* 102, 13325–13330. DOI: 10.1073/pnas.0505592102
- [27] Keep R. F., Jones H. C. (1990). Cortical microvessels during brain development: a morphometric study in the rat. *Microvasc. Res.* 40, 412–426. DOI: 10.1016/0026-2862(90)90036-Q
- [28] Khazipov R., Zaynutdinova D., Ogievetsky E., Valeeva G., Mitrukina O., Manent J.-B., et al. (2015). Atlas of the postnatal rat brain in stereotaxic coordinates. *Front. Neuroanat.* 9:161. DOI: 10.3389/fnana.2015.00161
- [29] Kirmse K., Witte O. W., Holthoff K. (2010). GABA depolarizes immature neocortical neurons in the presence of the Ketone body -Hydroxybutyrate. *J. Neurosci.* 30, 16002–16007. DOI: 10.1523/JNEUROSCI.2534-10.2010

- [30] Kocsis L., Herman P., Eke A. (2006). The modified BeerLambert law revisited. *Phys. Med. Biol.* 51, N91–N98. DOI: 10.1088/0031-9155/51/5/N02
- [31] Kohl M., Lindauer U., Royle G., Kuhl M., Gold L., Villringer A., et al. (2000). Physical model for the spectroscopic analysis of cortical intrinsic optical signals. *Phys. Med. Biol.* 45, 3749–3764. DOI: 10.1088/0031-9155/45/12/317
- [32] Kozberg M. G., Chen B. R., Deleo S. E., Bouchard M. B., Hillman E. M. C. (2013). Resolving the transition from negative to positive blood oxygen level-dependent responses in the developing brain. *Proc. Natl. Acad. Sci. U.S.A.* 110, 4380–4385. DOI: 10.1073/pnas.1212785110
- [33] Ma Y., Shaik M. A., Kim S. H., Kozberg M. G., Thibodeaux D. N., Zhao H. T., et al. (2016). Wide-field optical mapping of neural activity and brain haemodynamics: considerations and novel approaches. *Philos. Trans. R. Soc. Lond. Ser B Biol. Sci.* 371:20150360. DOI: 10.1098/rstb.2015.0360
- [34] Macvicar B. A., Feighan D., Brown A., Ransom B. (2002). Intrinsic optical signals in the rat optic nerve: role for K<sup>+</sup> uptake via NKCC1 and swelling of astrocytes. *Glia* 37, 114–123. DOI: 10.1002/glia.10023
- [35] MacVicar B. A., Hochman D. (1991). Imaging of synaptically evoked intrinsic optical signals in hippocampal slices. *J. Neurosci.* 11, 1458–1469.
- [36] Malonek D., Grinvald A. (1996). Interactions between electrical activity and cortical microcirculation revealed by imaging spectroscopy: implications for functional brain mapping. *Science* 272, 551–554.
- [37] Matcher S. J., Cope M., Delpy D. T. (1994). Use of the water absorption spectrum to quantify tissue chromophore concentration changes in near-infrared spectroscopy. *Phys. Med. Biol.* 39, 177–196. DOI: 10.1088/0031-9155/39/1/011
- [38] Mayevsky A., Chance B. (1982). Intracellular oxidation-reduction state measured in situ by a multichannel fiber-optic surface fluorometer. *Science* 217, 537–540.
- [39] Mayevsky A., Rogatsky G. G. (2007). Mitochondrial function in vivo evaluated by NADH fluorescence: from animal models to human studies. *Am. J. Physiol. Cell Physiol.* 292, 615–640. DOI: 10.1152/ajpcell.00249.2006
- [40] Minlebaev M., Ben-Ari Y., Khazipov R. (2007). Network mechanisms of spindle-burst oscillations in the neonatal rat barrel cortex in vivo. *J. Neurophysiol.* 97, 692–700. DOI: 10.1152/jn.00759.2006
- [41] Minlebaev M., Colonnese M., Tsintsadze T., Sirota A., Khazipov R. (2011). Early gamma oscillations synchronize developing thalamus and cortex. *Science* 334, 226–229. DOI: 10.1126/science.1210574
- [42] Mitrukina O., Suchkov D., Khazipov R., Minlebaev M. (2015). Imprecise whisker map in the neonatal rat barrel cortex. *Cereb. Cortex* 25, 3458–3467. DOI: 10.1093/cercor/bhu169
- [43] Norman M. G., O’Kusky J. R. (1986). The growth and development of microvasculature in human cerebral cortex. *J. Neuropathol. Exp. Neurol.* 45, 222–32.
- [44] Obrig H. (2014). NIRS in clinical neurology - a ‘promising’ tool? *Neuroimage* 85, 535–546. DOI: 10.1016/j.neuroimage.2013.03.045
- [45] Paxinos G., Watson C. (1997). *The Rat Brain in Stereotaxic Coordinates*, 3rd Edn. San Diego, CA: Academic Press.
- [46] Peterson B. E., Goldreich D. (1994). A new approach to optical imaging applied to rat barrel cortex. *J. Neurosci. Methods* 54, 39–47. DOI: 10.1016/0165-0270(94)90158-9
- [47] Rector D. M., Carter K. M., Volegov P. L., George J. S. (2005). Spatio-temporal mapping of rat whisker barrels with fast scattered light signals. *Neuroimage* 26, 619–627. DOI: 10.1016/j.neuroimage.2005.02.030
- [48] Risser L., Plouraboué F., Cloetens P., Fonta C. (2009). A 3D-investigation shows that angiogenesis in primate cerebral cortex mainly occurs at capillary level. *Int. J. Dev. Neurosci.* 27, 185–196. DOI: 10.1016/j.ijdevneu.2008.10.006
- [49] Rubin B. D., Katz L. C. (1999). Optical imaging of odorant representations in the mammalian olfactory bulb. *Neuron* 23, 499–511. DOI: 10.1016/S0896-6273(00)80803-X
- [50] Sheth S., Nemoto M., Guiou M., Walker M., Pouratian N., Toga A. W. (2003). Evaluation of coupling between optical intrinsic signals and neuronal activity in rat somatosensory cortex. *Neuroimage* 19, 884–894. DOI: 10.1016/S1053-8119(03)00086-7
- [51] Sheth S. A., Nemoto M., Guiou M., Walker M., Pouratian N., Toga A. W. (2004). Linear and nonlinear relationships between neuronal activity, oxygen metabolism, and hemodynamic responses. *Neuron* 42, 347–355. DOI: 10.1016/S0896-6273(04)00221-1
- [52] Sheth S. A., Nemoto M., Guiou M. W., Walker M. A., Toga A. W. (2005). Spatiotemporal evolution of functional hemodynamic changes and their relationship to neuronal activity. *J. Cereb. Blood Flow Metab.* 25, 830–841. DOI: 10.1038/sj.jcbfm.9600091
- [53] Strangman G., Culver J. P., Thompson J. H., Boas D. A. (2002). A quantitative comparison of simultaneous BOLD fMRI and NIRS recordings during functional brain activation. *Neuroimage* 17, 719–731. DOI: 10.1006/nimg.2002.1227
- [54] Valeeva G., Valiullina F., Khazipov R. (2013). Excitatory actions of GABA in the intact neonatal rodent hippocampus in vitro. *Front. Cell. Neurosci.* 7:20. DOI: 10.3389/fncel.2013.00020

- [55] Valiullina F., Akhmetshina D., Nasretudinov A., Mukhtarov M., Valeeva G., Khazipov R., et al. (2016). Developmental changes in electrophysiological properties and a transition from electrical to chemical coupling between excitatory layer 4 neurons in the rat barrel cortex. *Front. Neural Circ.* 10:1. DOI: 10.3389/fncir.2016.00001
- [56] van der Zee P., Essenpreis M., Delpy D. T. (1993). Optical properties of brain tissue, in *Photon Migration and Imaging in Random Media and Tissues*, Proc. SPIE Vol. 1888, eds Chance B., Alfano R. R., editors. (Bellingham, WA:), 454–465. DOI: 10.1117/12.154665
- [57] Villringer A., Planck J., Hock C., Schleinkofer L., Dirnagl U. (1993). Near infrared spectroscopy (NIRS): a new tool to study hemodynamic changes during activation of brain function in human adults. *Neurosci. Lett.* 154, 101–104.
- [58] Vincis R., Lagier S., Van De Ville D., Rodriguez I., Carleton A. (2015). Sensory-evoked intrinsic imaging signals in the olfactory bulb are independent of neurovascular coupling. *Cell Rep.* 12, 1–13. DOI: 10.1016/j.celrep.2015.06.016
- [59] Wang L. V., Jacques S. L., Zheng L. (1995). MCML - Monte Carlo modeling of light transport in multi-layered tissues. *Comput. Methods Prog. Biomed.* 47, 131–146.
- [60] Watanabe M., Fukuda A., Rivera C., Cherubini E. (2015). Development and regulation of chloride homeostasis in the central nervous system. *Front. Cell Neurosci.* 9:371. DOI: 10.3389/fncel.2015.00371
- [61] Workman A. D., Charvet C. J., Clancy B., Darlington R. B., Finlay B. L. (2013). Modeling transformations of neurodevelopmental sequences across mammalian species. *J. Neurosci.* 33, 7368–7383. DOI: 10.1523/JNEUROSCI.5746-12.2013
- [62] Wyatt J. S., Delpy D. T., Cope M., Wray S., Reynolds E. O. R. (1986). Quantification of cerebral oxygenation and haemodynamics in sick newborn infants by near infrared spectrophotometry. *Lancet* 328, 1063–1066.
- [63] Yamada H., Sadato N., Konishi Y., Muramoto S., Kimura K., Tanaka M., et al. (2000). A milestone for normal development of the infantile brain detected by functional MRI. *Neurology* 55, 218–223. DOI: 10.1212/WNL.55.2.218
- [64] Zehendner C. M., Tsohataridis S., Luhmann H. J., Yang J. W. (2013). Developmental switch in neurovascular coupling in the immature rodent barrel cortex. *PLoS ONE* 8:e80749. DOI: 10.1371/journal.pone.0080749
- [65] Zheng Y., Johnston D., Berwick J., Chen D., Billings S., Mayhew J. (2005). A three-compartment model of the hemodynamic response and oxygen delivery to brain. *Neuroimage* 28, 925–939. DOI: 10.1016/j.neuroimage.2005.06.042
- [66] Zijlstra W., Buursma A., Falke H., Catsburg J. (1994). Spectrophotometry of hemoglobin: absorption spectra of rat oxyhemoglobin, deoxyhemoglobin, carboxyhemoglobin, and methemoglobin. *Comp. Biochem. Physiol. B Comp. Biochem.* 107, 161–166. DOI: 10.1016/0305-0491(94)90238-0